# Neutron scattering research on organic solar cell materials

F.M. Mulder

O. Kruglova

L. van Eijck

G.J. Kearley

IRI, Delft University of Technology

S.J. Picken

J. Stride


B. Farago

TNW, Delft University of Technology

ILL, Grenoble, France




#### Group activities in energy storage and conversion materials





#### Solar energy applications

Team of Delft University wins World Solar Challenges 2001, 2003





#### Energy payback ratio =

usable energy delivered during lifetime

energy used for building & running device/plant

Solar cell (monocrystalline)	5 - 7
Solar cell (thin film, organic)	15 - 20
Wind	30 – 40
D-T Fusion	(27)
Fission	~ 16
Coal:	11 - 14
Gasoline	10 – 40 ??



#### Organic solar cell by self-assembly?


L. Schmidt-Mende et al. Science 293 (2001) 1119 & 1059

Grätzel type solar cells using organic dyes: efficiency up to 11%


Overview: see e.g. M. Grätzel, Nature 414 (2001) 338.



#### Hole conducting organic dye: Polythiophene










### Co-continuous network of perylene electron conductor and discotic hole conductor

Model compound for discotics: Liquid crystalline Between 69 and 99 °C hexa-alkoxy triphenylene: HAT6



#### Charge transport along columns:

- → inter-disc hopping on ps timescales
- → long-range transport over ~100 nm on ~ns



Could molecular motion influence charge transfer?

**ps** 

A.M. van de Craats et al. J. Phys. Chem. B102 (1998) 9625.



#### Quasi elastic neutron scattering (IN6, ILL, Grenoble)


4 ] \_\_\_\_ Observed

Unexpected: There is not much difference between the spectral shapes of

- fully protonated HAT6
- tail deuterated HAT6
- core deuterated HAT6


Energy Transfer (meV)








#### Force field molecular dynamics simulation




Out-of-plane motion on 7ps time-scale




Greater mobility in-plane on 0.2ps time-scale









June 4-5 2004, Washington



#### Force field molecular dynamics simulations suggest:


The experimentally observed motions could correspond to:


- Fast 7 ps motions: in plane amplitude: ~ 0.3 A
- Relatively slow 0.2 ps: out of plane motions amplitude: ~1.2A

F.M. Mulder, et al. J. Am. Chem. Soc. 2003, 125, 3860-3866




#### Can one observe anisotropic dynamics?









Change in Bragg-peaks in quasi-elastic spectra shows orientation worked

Orient by shear at ~365K, then keep > 350K





#### Result:

- not much difference between vertical and horizontal alignment
- both quasi elastic signals are still present




## Helical modes would have Similar amplitudes // and $\bot$ to column







#### Relevant scale for the rotation of HAT6 molecule?




$$\frac{2\pi}{d}$$
 ~ 0.2 Å<sup>-1</sup>



Neutron Spin Echo





#### NSE (IN15, ILL Grenoble)



Limit for rotational diffusion








 $Q = 0.2 \text{ Å}^{-1}$ 

 $Q = 0.1 \text{ Å}^{-1}$ 



#### Conductivity?



First step: influence of certain vibrational modes on electronic orbitals



#### Conclusions

- neutron scattering combined with simulations yields structure & dynamics
- molecular and electronic dynamics have similar timescales in HAT6
- simulations will yield conductivity, to be continued

'Molecular dynamics assisted' conductivity?



#### Acknowledgements



**Dutch Science Foundation** 




**Chemical Sciences Foundation** 



Delft Institute for Sustainable Energy















#### Slow dynamics of HAT6 observed by neutron spin echo

